

- Nardelli, M. (1983). *Comput. Chem.* **7**, 95–98.
 Pattenden, G. (1978). *Fortschr. Chem. Org. Naturst.* **35**, 133–136.
 Rao, Y. S. (1964). *Chem. Rev.* **64**, 353–383.
 Rao, Y. S. (1976). *Chem. Rev.* **76**, 625–682.
 Sheldrick, G. M. (1976). *SHELX76*. Program for crystal structure determination. Univ. of Cambridge, England.
 Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
 Věgh, D., Jakubcová, M., Kožisek, J., Bencková, M. & Dvorský, A. (1993). *Synth. Commun.* (In preparation.)
 Walker, N. & Stuart, D. (1983). *Acta Cryst.* **A39**, 158–166.

Acta Cryst. (1993). **C49**, 992–993

p-Bromobenzoate Derivative of 3-(*tert*-Butyldimethylsilyl)-5-hydroxy-2(5*H*)-furanone

MASOOD PARVEZ, SHUYUAN YU AND BRIAN A. KEAY

Department of Chemistry, The University of Calgary,
 2500 University Drive NW, Calgary, Alberta,
 Canada T2N 1N4


(Received 22 July 1992; accepted 3 November 1992)

Abstract

In 3-(*tert*-butyldimethylsilyl)-2-oxo-5*H*-5-furyl 4-bromobenzoate, the furan ring is almost planar with a maximum deviation of 0.047 (6) Å, while the benzene ring is essentially planar [maximum deviation 0.012 (5) Å]; the best mean planes of the two rings are inclined at an angle of 73.7°. The average bond distances are: Si—C 1.865 (7), C_{sp³}—C_{sp³} 1.523 (9), C_{sp²}—C_{sp²} 1.487 (7), C—C_{arom} 1.381 (7), C_{sp³}—O 1.423 (6), C_{sp²}—O 1.362 (6) and C=O 1.204 (6) Å.

Comment

Treatment of 2-(*tert*-butyldimethylsilyl)furan (1) with lead tetraacetate in acetic acid at 343 K for 2 h provided compounds (2) and (3) in a 2:1 ratio. The



major compound (2) was identified as 5-(*tert*-butyldimethylsilyl)-5-acetoxy-2(5*H*)-furanone by NMR, IR and MS spectra. The minor compound was identified by converting the acetate of compound (3) into a *p*-bromobenzoate (4) and performing an X-ray structure determination. The structure consists of discrete molecules separated by normal van der Waals distances.




Fig. 1. ORTEP (Johnson, 1976) drawing of the title compound showing the labelling of the non-H atoms. Thermal ellipsoids are shown at 50% probability levels; H atoms are assigned arbitrary radii.

Experimental

Crystal data

C₁₇H₂₁BrO₄Si
M_r = 397.34
 Triclinic
P $\bar{1}$
a = 7.350 (4) Å
b = 19.906 (10) Å
c = 6.726 (5) Å
 α = 93.93 (6)°
 β = 109.10 (6)°
 γ = 87.59 (5)°
V = 927 (1) Å³
Z = 2

D_x = 1.423 Mg m⁻³
 Mo *K*α radiation
 λ = 0.71069 Å
 Cell parameters from 25 reflections
 θ = 10–15°
 μ = 2.27 mm⁻¹
T = 152 (1) K
 Block
 0.50 × 0.40 × 0.30 mm
 Colourless

Data collection

Rigaku AFC-6S diffractometer
 $\omega/2\theta$ scans, width (1.78 + 0.35tan θ)°, rate 8.0° min⁻¹ in ω
 Absorption correction: empirical (North, Phillips & Mathews, 1968)
T_{min} = 0.723, *T_{max}* = 0.999
 3560 measured reflections

3279 independent reflections
 2289 observed reflections [*I* > 3σ(*I*)]
R_{int} = 0.0255
 θ_{max} = 25°
h = 0 → 8
k = -23 → 23
l = -7 → 7
 3 standard reflections
 frequency: 100 min
 intensity variation: <0.7%

Refinement

Refinement on *F*
 Final *R* = 0.048
 wR = 0.033
S = 3.49
 2289 reflections
 271 parameters

Only coordinates of H atoms refined
 $w = 1/(\sigma^2 F)$
 $(\Delta/\sigma)_{\text{max}}$ = 0.05
 $\Delta\rho_{\text{max}}$ = 0.68 e Å⁻³
 $\Delta\rho_{\text{min}}$ = -0.71 e Å⁻³

Table 1. Fractional atomic coordinates and equivalent isotropic thermal parameters (\AA^2)

$$B_{\text{eq}} = (8\pi^2/3)[(aa^*)^2 U_{11} + (bb^*)^2 U_{22} + (cc^*)^2 U_{33} + (2aa^*bb^*\cos\gamma)U_{12} + (2aa^*cc^*\cos\beta)U_{13} + (2bb^*cc^*\cos\alpha)U_{23}]$$

	x	y	z	B_{eq}
Br	-0.0747 (1)	0.3478 (1)	0.2192 (1)	0.0405 (4)
Si	0.2622 (2)	0.8764 (1)	0.2976 (2)	0.028 (1)
O(1)	0.6436 (5)	0.5627 (2)	0.2602 (5)	0.031 (2)
O(2)	0.4188 (5)	0.6344 (2)	0.3208 (5)	0.024 (2)
O(3)	0.5209 (5)	0.7036 (2)	0.1153 (5)	0.026 (2)
O(4)	0.3812 (5)	0.7850 (2)	-0.0984 (5)	0.035 (3)
C(1)	0.0930 (8)	0.4189 (3)	0.2263 (7)	0.025 (4)
C(2)	0.0399 (8)	0.4824 (3)	0.2661 (8)	0.024 (3)
C(3)	0.1642 (8)	0.5348 (3)	0.2815 (8)	0.023 (4)
C(4)	0.3447 (7)	0.5196 (3)	0.2575 (7)	0.020 (3)
C(5)	0.3925 (8)	0.4543 (3)	0.2127 (8)	0.026 (4)
C(6)	0.2681 (9)	0.4025 (3)	0.1948 (8)	0.029 (4)
C(7)	0.4865 (8)	0.5727 (3)	0.2780 (7)	0.024 (4)
C(8)	0.5384 (8)	0.6891 (3)	0.3255 (9)	0.030 (4)
C(9)	0.4639 (7)	0.7501 (3)	0.4129 (8)	0.023 (3)
C(10)	0.3936 (7)	0.7950 (3)	0.2696 (8)	0.024 (3)
C(11)	0.4241 (7)	0.7639 (3)	0.0747 (8)	0.022 (4)
C(12)	0.3853 (10)	0.9129 (3)	0.5655 (10)	0.035 (4)
C(13)	0.2723 (9)	0.9342 (3)	0.1005 (10)	0.039 (4)
C(14)	0.0030 (7)	0.8550 (3)	0.2581 (9)	0.035 (4)
C(15)	-0.1001 (10)	0.9171 (4)	0.3215 (13)	0.057 (6)
C(16)	-0.0978 (10)	0.8343 (4)	0.0286 (12)	0.061 (6)
C(17)	-0.0033 (10)	0.7982 (4)	0.3969 (13)	0.055 (5)

Table 2. Geometric parameters (\AA , $^\circ$)

Br—C(1)	1.904 (6)	Si—C(10)	1.881 (6)
Si—C(12)	1.845 (7)	Si—C(13)	1.836 (7)
Si—C(14)	1.898 (6)	O(1)—C(7)	1.206 (6)
O(2)—C(7)	1.355 (6)	O(2)—C(8)	1.420 (6)
O(3)—C(8)	1.426 (6)	O(3)—C(11)	1.368 (6)
O(4)—C(11)	1.202 (6)	C(1)—C(2)	1.343 (7)
C(1)—C(6)	1.393 (9)	C(2)—C(3)	1.391 (9)
C(3)—C(4)	1.406 (7)	C(4)—C(5)	1.372 (7)
C(4)—C(7)	1.484 (7)	C(5)—C(6)	1.380 (8)
C(8)—C(9)	1.476 (7)	C(9)—C(10)	1.321 (7)
C(10)—C(11)	1.490 (7)	C(14)—C(15)	1.531 (9)
C(14)—C(16)	1.513 (9)	C(14)—C(17)	1.526 (11)
C(10)—Si—C(12)	106.4 (3)	C(10)—Si—C(13)	110.5 (3)
C(10)—Si—C(14)	107.2 (3)	C(12)—Si—C(13)	110.1 (3)
C(12)—Si—C(14)	112.1 (3)	C(13)—Si—C(14)	110.3 (3)
C(7)—O(2)—C(8)	115.6 (4)	C(8)—O(3)—C(11)	107.6 (4)
Br—C(1)—C(2)	118.8 (5)	Br—C(1)—C(6)	118.4 (5)
C(2)—C(1)—C(6)	122.8 (6)	C(1)—C(2)—C(3)	119.7 (6)
C(2)—C(3)—C(4)	118.8 (5)	C(3)—C(4)—C(5)	119.9 (5)
C(3)—C(4)—C(7)	121.7 (5)	C(5)—C(4)—C(7)	118.4 (5)
C(4)—C(5)—C(6)	121.2 (5)	C(1)—C(6)—C(5)	117.5 (6)
O(1)—C(7)—O(2)	123.5 (6)	O(1)—C(7)—C(4)	124.8 (6)
O(2)—C(7)—C(4)	111.7 (5)	O(2)—C(8)—O(3)	109.4 (5)
O(2)—C(8)—C(9)	108.6 (5)	O(3)—C(8)—C(9)	105.1 (5)
C(8)—C(9)—C(10)	111.8 (5)	Si—C(10)—C(9)	127.5 (4)
Si—C(10)—C(11)	127.3 (4)	C(9)—C(10)—C(11)	104.9 (5)
O(3)—C(11)—O(4)	120.0 (5)	O(3)—C(11)—C(10)	110.2 (5)
O(4)—C(11)—C(10)	129.8 (5)	Si—C(14)—C(15)	109.2 (5)
Si—C(14)—C(16)	109.6 (5)	Si—C(14)—C(17)	110.1 (4)
C(15)—C(14)—C(16)	109.3 (6)	C(15)—C(14)—C(17)	108.4 (6)
C(16)—C(14)—C(17)	110.2 (6)		

Space group $P\bar{1}$ or $P1$; the former was assumed and confirmed by successful analysis. Lorentz-polarization corrections were applied but not extinction corrections. The structure was solved by the heavy-atom method and refined by full-matrix least squares with the non-H atoms anisotropic. The H atoms were located from the ΔF map and allowed to refine with fixed isotropic temperature factors. The scattering factors were taken from Cromer & Waber (1974) and Stewart, Davidson & Simpson (1965); allowance was made for anomalous dispersion (Ibers & Hamilton, 1964). All calculations were performed using TEXSAN

(Molecular Structure Corporation, 1992) on a Silicongraphics Personal Iris D/35 computer. A search of the Cambridge Structural Database (Allen, Kennard & Taylor, 1983) yielded only one hit of a similar compound, a diphenylmethylsilylbenzofuran (de Perez, Fuentes, Larson, Barnes & Heeg, 1986).

Financial support from the Natural Sciences and Engineering Research Council of Canada and the University of Calgary Research Board is gratefully acknowledged.

Lists of structure factors, anisotropic thermal parameters, H-atom coordinates and complete geometry have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 55823 (22 pp.). Copies may be obtained through The Technical Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England. [CIF reference: BR1021]

References

- Allen, F. H., Kennard, O. & Taylor, R. (1983). *Acc. Chem. Res.* **16**, 146–153.
- Cromer, D. T. & Waber, J. T. (1974). *International Tables for X-ray Crystallography*, Vol. IV, Table 2.2A. Birmingham: Kynoch Press. (Present distributor Kluwer Academic Publishers, Dordrecht.)
- Ibers, J. A. & Hamilton, W. C. (1964). *Acta Cryst.* **17**, 781–782.
- Johnson, C. K. (1976). *ORTEP*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Molecular Structure Corporation (1992). *TEXSAN*. Version 1.2. Single-crystal structure analysis software. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Perez, R. M. B. de, Fuentes, L. M., Larson, G. L., Barnes, C. L. & Heeg, M. J. (1986). *J. Org. Chem.* **51**, 2039–2043.
- Stewart, R. F., Davidson, E. R. & Simpson, W. T. (1965). *J. Chem. Phys.* **42**, 3175–3187.

Acta Cryst. (1993). **C49**, 993–996

Structure of Methyl 2-(Nitrooxy)ethyl 1,4-Dihydro-2,6-dimethyl-4-(3-nitrophenyl)-3,5-pyridinedicarboxylate

T. OGAWA, K. MATSUMOTO,* C. YOKOO,
K. HATAYAMA AND K. KITAMURA

Research Center, Taisho Pharmaceutical Co., Ltd,
1-403 Yoshino-cho, Ohmiya 330, Japan

(Received 10 August 1992; accepted 24 November 1992)

Abstract

The orientations of the carbonyl groups at C3 and C5 are different. The phenyl ring linked to C4 is perpendicular to the dihydropyridine ring. Some other structural features have also been elucidated.